

Winterthurerstr. 190 CH-8057 Zurich http://www.zora.uzh.ch

Voar.	2	0	n	Q
1 P.U.I	7.1	•	,	()

# Contour copying or echoplasia - a new echo phenomenon in a person with Gilles de la Tourette syndrome

Kawohl, W; Podoll, K

Kawohl, W; Podoll, K (2008). Contour copying or echoplasia - a new echo phenomenon in a person with Gilles de la Tourette syndrome. Psychopathology, 41(3):201-202.

Postprint available at: http://www.zora.uzh.ch

Posted at the Zurich Open Repository and Archive, University of Zurich. http://www.zora.uzh.ch

Originally published at: Psychopathology 2008, 41(3):201-202.

### Letter to the Editor

### **Psychopathology**

Psychopathology 2008;41:201–202 DOI: 10.1159/000120989 Received: July 2, 2007 Accepted: July 12, 2007 Published online: March 13, 2008

## Contour Copying or Echoplasia – A New Echo Phenomenon in a Person with Gilles de la Tourette Syndrome

W. Kawohl<sup>a</sup> K. Podoll<sup>b</sup>

<sup>a</sup>Research Group Clinical and Experimental Psychopathology, Psychiatric University Hospital Zurich, Zurich, Switzerland; <sup>b</sup>Department of Psychiatry, RWTH Aachen University, Aachen, Germany

Echo phenomena are environment-driven responses [1] occurring in different disorders, e.g. the Gilles de la Tourette syndrome (GTS) [2, 3], schizophrenic disorders [4, 5] including catatonia [6, 7], aphasias [8, 9], autism [10–12], epilepsy [13], frontotemporal dementia [14], Pick's disease [15] and neuropsychiatric systemic lupus erythematosus [16]. They are defined as pathological imitations of external stimuli. We report the case of a patient with GTS presenting a, to our knowledge, yet unpublished echo phenomenon.

A 43-year-old, right-handed father of a family was sent to our outpatient clinic by his general practitioner for having symptoms interpreted as characteristic of attention-deficit hyperactivity syndrome. During the exploration he spoke in a loud way and presented different facial tics. He also reported different single vocal tics, the naming of objects and echolalia. His father had similar symptoms, albeit more pronounced, as well as impulsive outbreaks. His 16-year-old daughter had minor symptoms of the same quality. Being asked for more phenomena similar to the echolalia, he reported that when seeing something impressive such as a big building, an attractive woman or a sports car he would draw the shape of the object with his right index finger, either in the air or on any surface within reach (e.g. on his own clothing). Whether he would

draw in the air or on an available surface (e.g. on his leg) depended on the position of his hand at the moment of perceiving the stimulus. Sometimes he would only trace the contours in his mind. Any premonitory urge was denied. If suppressed, a feeling of pressure, but not of fear, would occur.

Physical examination was normal. The patient refused any medication for the treatment of his tics, denying any psychological strain. The patient met the criteria for the diagnosis of GTS according to DSM-IV as well as to ICD-10. He neither fulfilled the criteria for a co-morbid obsessive-compulsive disorder nor for an attention-deficit hyperactivity syndrome.

Despite a considerable amount of literature on repetitive phenomena, the symptom reported here has, as far as we know, not been described and categorized before. There are some superficial similarities to obsessive-compulsive symptoms, yet the symptom presented by our patient does not belong to them. When obsessive acts are suppressed, apprehensions that something bad might happen occur frequently. These apprehensions do not occur in the case of tics [2]. In contrast to compulsions, neither the act itself nor its suppression causes any fear. Cath et al. [17] emphasize that symptoms in GTS, even when sharing superficial similarities with obsessive-compulsive symptoms,

should not be diagnosed automatically as obsessive compulsive. This can also be applied to the phenomenon reported here: unlike, for instance, obsessive drawing on a table cloth in an obsessive-compulsive disorder patient, our patient's tic occurs with an external model as a stimulus. In summary, the characteristics of this particular symptom correspond to the definition of an echo phenomenon.

Echo phenomena are classified as tics [18]. In GTS, echo phenomena are associated with illness duration and the severity of motor tics [3]. Echo phenomena reported in the literature are echolalia, the repetition of words, echokinesis or echopraxia, the reproduction of movements, echomimia, the reproduction of facial expressions, and echographia, the repetition of written [19] and spoken words [20]. Echo phenomena are named by merging the Greek word 'echo' and the word that describes the repeated action, e.g. echolalia consisting of 'echo' and 'lalein' (to speak). To distinguish this phenomenon from other echo phenomena we propose the terminus 'echoplasia'. Echoplasia is, in analogy to the terminology of the other echo phenomena, a compound word composed of 'echo' and 'plathein', the Greek word for 'to shape'. We suggest that this phenomenon should be added to the list of echo phe-

### **Acknowledgements**

The authors would like to thank Dr. phil. Th. Optenbrenk, Nettetal, Dr. phil. I Kawohl, Mönchengladbach, Germany, and lic. phil. Hartmann, University of Zurich, Switzerland, for their terminological help including the verification of the Greek translations. We also wish to thank David Feder, M.Sc. (Psychiatric University Hospital, Zurich) for proof reading the manuscript.

#### References

- 1 Ghika J, Tennis M, Growdon J, Hoffman E, Johnson K: Environment-driven responses in progressive supranuclear palsy. J Neurol Sci 1995;130:104–111.
- 2 Müller N: Tics; in Jahn T (ed): Bewegungsstörungenbeipsychischen Erkrankungen. Berlin, Springer, 2004, pp 215–231.
- 3 Robertson MM, Trimble MR, Lees AJ: The psychopathology of the Gilles de la Tourette syndrome. A phenomenological analysis. Br J Psychiatry 1988;152:383–390.
- 4 Kirov K: Long-term study of schizophasic patients. Psychopathology 1985;18:226–229.

- 5 Kirov K: Studies of the course of schizophasia. Psychopathology 1990;23:339–341.
- 6 Ford RA: The psychopathology of echophenomena. Psychol Med 1989;19:627–635.
- 7 Lee JW: Chronic 'speech catatonia' with constant logorrhea, verbigeration and echolalia successfully treated with lorazepam: a case report. Psychiatry Clin Neurosci 2004;58: 666–668.
- 8 Ghika J, Bogousslavsky J, Ghika-Schmid F, Regli F: 'Echoing approval': a new speech disorder. J Neurol 1996;243:633–637.
- 9 Christman SS, Boutsen FR, Buckingham HW: Perseveration and other repetitive verbal behaviors: functional dissociations. Semin Speech Lang 2004;25:295–307.
- 10 Williams JH, Whiten A, Suddendorf T, Perrett DI: Imitation, mirror neurons and autism. Neurosci Biobehav Rev 2001;25:287– 295.
- 11 Schuler AL: Beyond echoplaylia: promoting language in children with autism. Autism 2003;7:455–469.
- 12 Prizant BM, Duchan JF: The functions of immediate echolalia in autistic children. J Speech Hear Disord 1981;46:241–249.
- 13 Linetsky E, Planer D, Ben-Hur T: Echolaliapalilalia as the sole manifestation of nonconvulsive status epilepticus. Neurology 2000; 55:733-734.

- 14 Bathgate D, Snowden JS, Varma A, Blackshaw A, Neary D: Behaviour in frontotemporal dementia, Alzheimer's disease and vascular dementia. Acta Neurol Scand 2001;103: 367–378.
- 15 Mimura M, Oda T, Tsuchiya K, Kato M, Ikeda K, Hori K, Kashima H: Corticobasal degeneration presenting with nonfluent primary progressive aphasia: a clinicopathological study. J Neurol Sci 2001;183:19–26.
- 16 Zapor M, Murphy FT, Enzenauer R: Echolalia as a novel manifestation of neuropsychiatric systemic lupus erythematosus. South Med J 2001;94:70–72.
- 17 Cath DC, van de Wetering BJ, van Woerkom TC, Hoogduin CA, Roos RA, Rooijmans HG: Mental play in Gilles de la Tourette's syndrome and obsessive-compulsive disorder. Br J Psychiatry 1992;161:542–545.
- 18 Shapiro T, Lucy P: Echoing in autistic children: a chronometric study of semantic processing. J Child Psychol Psychiatry 1978;19: 373–378.
- 19 Peters UH: Wörterbuch der Psychiatrie und medizinischen Psychologie. Munich, Urban und Schwarzenberg, 1990.
- 20 Kan R, Takahashi Y, Takahashi R, Hagiwara M, Yashima Y, Kumashiro H: Epileptic patient demonstrating echographia during interictal states. Jpn J Psychiatry Neurol 1990; 44:404–406.